

MĂGURA - NEW VARIETY OF VINE FOR RED WINES CREATED AT S.C.D.V.V. ODOBEȘTI

MĂGURA - SOI NOU PENTRU OBTINEREA VINURILOR ROȘII DE CALITATE CREAT LA S.C.D.V.V. ODOBEȘTI

BOSOI Marioara¹, MIHU G. ¹, BOSOI Ionica¹, STOIAN Ileana¹
e-mail: mioara_bosoi@yahoo.com

Abstract. *The studying the long-term of vine germplasm, results in choosing the most valuable genitors and perform a variety of intra- and interspecific hybridisation with obtaining valuable new genotypes, tolerant to pests and diseases and resistance to stressors. Of the many elite hybrid obtained at S.C.D.V.V. Odobești, in 2014 was approved the elite hybrid 18-46 under the name 'Măgura' - variety for quality red wines obtained by hybridization sexual of variety Babească neagră with the hybrid combination (Merlot x Alicante Bouschet). The new creation is characterized by middle-sized grapes (175g), small to medium berry (3.3 g), colored in black-blue, rich in anthocyanins in both the skin and core. The average grape production is 4.5 kg/vine or 16.6 tons/ha. It has a good resistance to the main cryptogamic diseases. The grapes reach maturity in the epoch IV. The wines obtained are extractive, intensely colored, tinctorial.*

Key words: variety, hybridization, elite hybrid, biological resistance

Rezumat. *Studierea pe termen lung a germoplasmei viticole are ca rezultat alegerea celor mai valoroși genitori și efectuarea unei game largi de hibridări intra și interspecifice cu obținerea de genotipuri noi valoroase, cu toleranță la boli și rezistențe la factorii de stres. Dintre numeroasele elite hibride obținute la S.C.D.V.V. Odobești, în anul 2014 a fost omologată elita hibridă 18-46 sub denumirea 'Măgura' - soi pentru vinuri roșii de calitate, obținut prin hibridarea sexuală a soiului Babească neagră cu combinația hibridă (Merlot x Alicante Bouschet). Noua creație se caracterizează prin struguri de mărime mijlocie (175 g), boabe mici spre mijlocii (3,3 g), colorate în negru - albastrui, bogate în antociani, atât în pielea cât și în miez. Producția medie de struguri este de 4,5 kg/butuc, respectiv 16,6 tone/ha. Are rezistență bună la principalele boli criptogamice. Strugurii ajung la maturitate în epoca a IV-a. Vinurile obținute sunt extractive, intens colorate, tinctoriale.*

Cuvinte cheie: soi, hibridare sexuală, elită hibridă, rezistență biologică

INTRODUCTION

The value of the varieties for obtaining the red wines is show both to level and the constant of grape production, its quality, defined by the ability of accumulation of sugars and anthocyanins in beans, as well as their tolerance to diseases and stress conditions. Research conducted in the last four decades in

¹ Research and Development Station for Viticulture and Oenology Odobești, Romania

Romania, led to valuable vine, with high tolerance to disease, resistant to drought and frost (Moldovan *et al.*, 1994, Oprea *et al.*, 2007 Calistru and Damian 1999, Culcea *et al.*, 2004).

The valorization of valuable hybrid combinations with genetic resistance, obtained over time in scientific research in the area of improving the vine, is one of the major objectives of Research and Development Station for Viticulture and Oenology Odobești. In this context it has created a new variety called "Măgura", which completes the varietal range of the varieties for obtain red wines quality, adapted to climatic conditions in vineyards in southeastern of Moldova.

MATERIAL AND METHOD

The study was carried out between 2012 - 2013, on a plantation aged 25 years, planting on soil type levigated,, located in the biological field SCDVV Odobești. For comparison, as a witness was used Băbească neagră variety, that represents the parents maternal and is similar in direction of production and epidermis' color.

The varieties Măgura and Băbească neagră was grafted on Kobber 5 BB rootstock, the training system with trunk of hub to the ground, with Dr. Guyot cutting system. Fruit load was 44 eye/hub, distributed on the canes with 9 eye and the spur with 2 eye. Distance of planting by 2.2 m x 1.2 m is returning 3788 but/ha. Were studied the ampelographic main characters, were made measurements and determinations on elements of fertility and productivity, the amount and quality of grape production, disease resistance, physico-chemical characteristics of the wine. The main climatic conditions of during the study period and multiannual values are presented in table 1.

Table 1

The main climatic conditions of study period (Odobești, 2012-2013)

| Climatic Indicator | Multiannual (1946 -2011) | Crop year | | Average 2012 - 2013 |
|---|-----------------------------|------------|------------|---------------------------|
| | | 2011 -2012 | 2012 -2013 | |
| Annual | | | | |
| The average temp., °C | 10.5 | 11.8 | 11.2 | 11,. |
| Temp max. abs. °C | 39.4 | 39.2 | 36.8 | 38.0 |
| Temp min. abs., °C | -22.8 | -22.4 | -17.3 | -19.9 |
| Amount degrees usuful temp.(Σ°tu) °C | 1604.0 | 1961.4 | 1879.2 | 1920.2 |
| The amount heatstroke hours | 2113.9 | 2585.1 | 2318.9 | 2452.0 |
| Precipitation amount, mm | 617.3 | 382.4 | 800.8 | 591.6 |
| On the vegetation period | | | | |
| The average temp., °C | 16.9 | 19.7 | 18.4 | 19.0 |
| Temp max. abs. °C | 39.4 | 39.2 | 36.8 | 38.0 |
| Temp min. abs., °C | -8.2 | -0.3 | -7.5 | -5.8 |
| Amount degrees usuful temp.(Σ°tu) °C | 1581.2 | 1938.0 | 1865.0 | 1901.5 |
| The amount heatstroke hours | 1629.7 | 1949.8 | 1842.9 | 1896.4 |
| Precipitation amount, mm | 431.2 | 327.2 | 561.8 | 444.5 |

RESULTS AND DISCUSSIONS

Research period was characterized by availability heliothermic large, but with very low rainfall in 2012 and very high in 2013, unevenly distributed during the growing season. Amount degrees useful temperature was 1920.2 compared to multiannual value of 1604.0 and the amount of annual rainfall was 591.6 mm, of which 444.5 mm during the growing season compared to multi annual value (617.3 mm), of which 431.2 mm during the growing season.

On the background of this weather conditions, Măgura variety has started its vegetation through disbudding between 19th - 23th of April, without significant differences in comparison with the witness. The flowering phenophase located between 23th-25th of May, and the veraison recorded between 25th - 30th of July. Grapes' ripening was realized between 25th of August - 5th of September, with approximately two weeks earlier than the witness variety (tab. 2).

In regards as the phenological spectrum cover in Odobești ecosystem conditions, the new variety Măgura completed their vegetation period after 190 days. Is the variety with the half-erlier maturation, grapes reach to full maturity in the first decade of September (epoch IV -a).

Table 2

Phenological spectrum in the conditions of Odobești vineyard

| Variety | Disbudding | Flowering | Veraison | Physiological maturity | Fall leaves |
|-----------------------|------------|-----------|------------|------------------------|-------------|
| Măgura | 19-23.04 | 22-25.05 | 25-30.07 | 25.08 -05.09 | 31.10-10.11 |
| Băbească neagră (Mt.) | 20-25.04 | 23-27.05 | 05 - 08.08 | 18.09- 18.09 | 04.11-15.11 |

Fertility and productivity of the new variety appreciated by the percentage of fertile shoots, fertility coefficients (absolute and relative) and productivity indices (absolute and relative) shows the superiority of the new creation for the elements analyzed in comparison with to the witness. The primary buds' viability had values close to witness (tab. 3).

Table 3

Viability and the fertility/ productivity elements of the Măgura variety
(Odobești, average data 2012 – 2013)

| Variety | Viability (%) | Fertile shoots (%) | Fertility coefficients | | Productivity indices | |
|-----------------------|---------------|--------------------|------------------------|----------|----------------------|----------|
| | | | Relative | Absolute | Relative | Absolute |
| Măgura | 89.12 | 72.0 | 1.04 | 1.44 | 197.6 | 274.4 |
| Băbească neagră (Mt.) | 91.30 | 63.3 | 0.77 | 1.21 | 168.7 | 266.3 |

The Măgura variety show a potential fertility of 72.0%, higher than the control (63.3%). The same hierarchy exists and in the case the absolute fertility coefficient (1.44 to Măgura variety, respectively 1.21 to Băbească neagră) and the absolute productivity index (274.4 – Măgura, respectively 266.3 - Băbească neagră).

The main characters ampelographic of Măgura variety (fig. 1 and fig. 2), according 2nd edition of the OIV Descriptor list for grape varieties and *Vitis* species, 2009.

- Young shoot
 - Opening of the shoot tip: half open;
 - Intensity of anthocyanin coloration: low;
 - Density of prostrate hairs on the shoot tip: low;
- Young leaf
 - Upper side color of blade: green areas anthocyanins;
- Shoot:
 - Attitude (before tying); semi-erect
 - Color of the dorsal side of internodes: green with red;
 - Color of the ventral side of internodes: green with red;
 - Length of tendrils: medium/long;
- Mature leaf
 - Size of blade: small;
 - Shape of blade: pentagonal;
 - Number of lobes: five;
 - Shape of teeth: both sides straight;
 - Degree of opening/overlapping of petiole sinus: overlapped;
 - Depth of upper lateral sinuses: deep;
 - Density of erect hairs on main veins on lower side of blade: medium;
 - length of petiole compared to length of middle vein: slightly shorter;



Fig. 1 Young shoot




Fig. 2 Măgura variety (adult leaf, grape)

- Bunch - length (peduncle excluded): medium;
 - width: medium;
 - Density: middle;
 - Peduncle length: short;
- Berry
 - Length: short;
 - Width: narrow
 - Shape: globose;

- color of skin: blue black;
- thickness of skin: thick;
- intensity of flesh anthocyanin coloration: very strong;
- firmness of flesh: slightly firm;

Studying the technological characteristics of grape production was completed knowledge elements for the new grape variety (tab. 4).

Table 4

The technological properties of the Măgura variety

(Odobesti, average data 2012 – 2013)

| Variety | No. bunch/vine | Average weight grapes (g) | Average weight 100 berry (g) | Production grape | | Sugars g/l | Total acidity g/l H ₂ SO ₄ |
|-----------------------|----------------|---------------------------|------------------------------|------------------|------|------------|--|
| | | | | kg./vine | t/ha | | |
| Măgura | 24.0 | 190.0 | 185.0 | 4.5 | 16.6 | 212.0 | 4.2 |
| Băbească neagră (Mt.) | 19.1 | 220.0 | 215.0 | 4.2 | 15.9 | 208.0 | 5.3 |

Though the average weight of bunch at Măgura variety is smaller (190 g) than the reference variety (220 g), the higher number of grapes per vine to determined a higher production (4.5 kg/vine) compared to a reference variety - Băbească neagră (4.2 kg/vine). As regards the quality of grape production, sugar content in the juice at the variety Măgura was 212.0 g/L (with 4 g higher than the control) and total acidity of 4.2 g/L H₂SO₄ (lower 1.1 g/L H₂SO₄ than the control).

In the climatic conditions of the years 2012 - 2013, the Magura variety manifested high resistance to the main cryptogamic diseases compared to the control – Băbească neagră (tab. 5).

Table 5

The behavior at the main diseases of the vine

(OIV descriptor list for grape varieties and *Vitis* species, 2nd edition – 2009)

| Variety | Downy mildew (<i>Plasmopara viticola</i>) | | Powdery mildew (<i>Uncinula necator</i>) | | Black rot (<i>Botrytis cinerea</i>) | |
|-----------------------|--|---------------|---|---------------|--|---------------|
| | Leaf OIV 452 | Grape OIV 453 | Leaf OIV 455 | Grape OIV 456 | Leaf OIV 458 | Grape OIV 459 |
| Măgura | 7 - 9 | 7 - 9 | 7 - 9 | 7 - 9 | 7 | 5 - 7 |
| Băbească neagră (Mt.) | 5 - 7 | 5 | 5 | 5 | 7 | 5 |

Main characteristics of the wines produced are shown in table 6.

Table 6

Physico-chemical characteristics of wines (average data 2012 – 2013)

| Variety | Alcohol vol. % | Total acidity g/l H ₂ SO ₄ | Dry extract unreducible g/l | Anthocyanins mg/l |
|-----------------------|----------------|--|-----------------------------|-------------------|
| Măgura | 13.38 | 4.83 | 22.24 | 442.8 |
| Băbească neagră (Mt.) | 13.25 | 5.31 | 18.49 | 358.7 |

The wine made from the variety Măgura had an alcoholic strength of 13.38% vol., upper the witness (Băbească neagră) with 0.13% vol. Total acidity was 4.83 g/L H₂SO₄, less with 0.48 g/L H₂SO₄ compared to the control (5.31 g/L H₂SO₄). Also anthocyanin content in wine (442.8 mg/L) is superior than the witness (358.7 mg/L). Higher values were obtained and for unreducible dry extract (3.75 mg/L greater than the witness).

The values of the technological indices resulted from the mechanical analysis of the grapes, complete the properties of the Măgura variety (tab. 7).

Table 7

Mechanical composition of grapes for Măgura variety (average data 2012 – 2013)

| Elements determined | | Măgura | Băbească neagră (Mt.) |
|------------------------|---------------------------------|--------|-----------------------|
| 1 kg grapes: | berry, g | 959.5 | 960.5 |
| | bunch, g | 40.5 | 39.5 |
| | no. berry normally developed | 530 | 445 |
| | must, g | 796 | 780 |
| | volume of must, cm ³ | 657 | 635 |
| 100 berry: | average weight, g | 185 | 215 |
| | skin weight, g | 21 | 32.5 |
| | core weight, g | 157.8 | 175 |
| | seeds weight, g | 6.2 | 7.5 |
| | number of seeds | 165 | 167 |
| | weight of 100 seeds, g | 3.7 | 3.3 |
| Technological indices: | berry index | 53 | 45 |
| | structure of the grape index | 23.69 | 24.31 |
| | composition of berry index | 7.51 | 5.38 |

CONCLUSIONS

1. Variety Măgura stems from a hybridization of Băbească neagră variety with the hybrid combination (Merlot x Alicante Bouschet). During the study period it maintained qualities of distinctness, uniformity and stability.

2. In comparison with the control (Băbească neagră), the elite has shown potential for high productivity and constant throughout the study period.

3. The elite has proved superior than the witness in terms of grape production / vine and grape sugar content;

4. Being a variety with genetic tolerance is recommended introduction into the vineyards suitable for green technologies.

REFERENCES

1. Calistru Gh., Damian Doina, 1999 – *Golia, soi nou de viță de vie cu rezistență sporită la ger*. Lucr. st. seria Horticultura, vol nr. 42, USAMV Iași.
2. Culcea Valeria, Preda Daniela, Marian I., 2004 – *Rosina, soi rezistent pentru vinuri albe*. Analele ICDVV, vol. XVII, București, pp.61-65.
3. Oprea St., Moldovan S.D., 2007 – *Ameliorarea viței de vie în România*. Editura Poliam, Cluj Napoca.
4. ***, 2009 – The 2nd edition of the OIV Descriptor list for grape varieties and Vitis species